

Isolation and analysis of the molecular epidemiology and zoonotic significance of *Mycobacterium tuberculosis* in domestic and wildlife ruminants from three states in India

This paper (No. 19112018-00136-EN) has been peer-reviewed, accepted, edited, and corrected by authors. It has not yet been formatted for printing. It will be published in December 2018 in issue 37 (3) of the *Scientific and Technical Review*

F. Mukherjee ^(1, a), V.S. Bahekar ^(1, 2, a), S.Y. Pasha ⁽³⁾, P. Kannan ⁽⁴⁾, A. Prasad ⁽¹⁾, S.K. Rana ⁽¹⁾, A. Kanani ⁽⁵⁾, G.K. Sharma ⁽⁶⁾, D. Premalatha ⁽⁷⁾ & V.A. Srinivasan ^{(8)*}

(1) Research and Development Laboratory, National Dairy Development Board, Gachibowli, Hyderabad 500032, Telangana, India

(2) Department of Biotechnology, Jawaharlal Nehru Technological University, Hyderabad 500085, Telangana, India

(3) Research and Development Laboratory, Indian Immunologicals Limited, Gachibowli, Hyderabad 500032, Telangana, India

(4) Department of Immunology, National Institute for Research in Tuberculosis, Chennai 600031, Tamil Nadu, India

(5) Department of Animal Husbandry, Government of Gujarat, Gandhinagar, Gujarat, India

(6) National, Dairy Development Board, Anand 38801, Gujarat, India

(7) Department of Biotechnology, St Martin's Engineering College, Hyderabad, India

(8) National, Dairy Development Board, 33 Telecom Nagar, Gachibowli, Hyderabad 500032, Telangana, India

(a) Authors contributed equally

*Corresponding author: srinivasanva1948@gmail.com

Summary

The majority of tuberculosis cases in ruminants are caused by *Mycobacterium bovis* (MB). However, in this study, the authors reported the isolation of *Mycobacterium tuberculosis* (MT) from bovine milk, nasal swabs and post-mortem tissue samples ($n = 841$) collected from cattle and buffaloes in the states of Telangana, Maharashtra and Gujarat in India in the period from 2010 to 2015. The isolates ($n = 7$) were confirmed as *Mycobacterium* due to their growth characteristics and colony morphology in a commercial liquid medium Mycobacteria Growth Indicator Tube (MGIT)TM employing the BD BACTECTM MGITTM 960 system and the Lowenstein Jensen (LJ) medium supplemented with glycerol but not with sodium pyruvate, and BD-DIFCOTM Middlebrook 7H10 agar containing oleic albumin dextrose catalase (OADC). These isolates were initially identified as members of the *Mycobacterium tuberculosis* complex (MTC) using a commercial nested polymerase chain reaction (PCR) kit based on the IS6110 MTC specific nucleotide sequence. The isolates were confirmed as MT using three commercial line probe assay kits, were further genotyped, and the spoligotypes identified were of East African Indian (EAI) 3_IND, EAI5, Central-Asian (CAS) 1_DELHI, U and T1 lineages. Two MT isolates from one antelope (*Antelope cervipara*) and one gazelle (*Gazelle bennettii*) from Gujarat, which were identified previously, were spoligotyped during this study and identified as belonging to EAI3_IND and EAI5 lineages, respectively. The epidemiological significance and zoonotic implications of regional presence and documentation of the same or two different spoligotypes in different species within the family *Bovidae* as well as humans is discussed.

Keywords

Epidemiology – India – Isolation – *Mycobacterium tuberculosis* – Ruminant – Spoligotyping– Zoonosis.

Introduction

Tuberculosis is a contagious chronic emaciating zoonotic disease that can infect a large number of animal species, mainly mammals, including domestic and wild ruminants, marine mammals and humans (1). Bovine tuberculosis (BT) has been reported as the causative agent for significant economic losses in the dairy industry in Turkey, Ethiopia and Argentina, ranging from US\$ 15 to 63 million (2, 3, 4). The disease is mostly caused by infection with a member of the *Mycobacterium tuberculosis* complex (MTC), namely *Mycobacterium bovis* (MB), while another member of the MTC, *Mycobacterium tuberculosis* (MT), is generally responsible for human tuberculosis (1, 3, 5). However, during the past decade, bovines infected with MT have been reported in Ethiopia, Turkey, the People's Republic of China, Nigeria and Spain (6, 7, 8, 9, 10), and also, in India (11, 12). In addition, in the district of Gandhinagar, Gujarat state, in western India, captive wild ruminants infected with MT have been reported (13). Moreover, the concurrent infection of the bovine population in India with MT and MB has also been reported (12, 14, 15, 16).

Genotyping of disease strains leads to the discovery of epidemiologically significant information and it is therefore considered an important tool for decision-making with respect to tuberculosis control programmes. Spacer oligonucleotide typing (spoligotyping), based on the analysis of a direct repeat (DR) locus, is one such globally recognised genotyping method (17). Although spoligotyping data on MT strains of human origin isolated from various geographical regions in India are available, similar information on MT isolates from domestic and wild ruminants from diverse regions of the country has yet to be recorded. In those studies conducted in India to date, only a single report on the genotyping of MT isolates from cattle, from the state of Himachal Pradesh, is available (12). This paper reported on the isolation of MT in 2006 from a lung lesion taken from a Holstein-Friesian cow, and from the broncho-mediastinal lymph node of a Jersey cow. Both animals were 14 years old and were from the district of Palampur in the state of

Himachal Pradesh, northern India. The spoligotype patterns on the X-ray film for the two isolates cited above, were further interpreted by the authors at the National Institute for Research in Tuberculosis (NIRT), Chennai, India, and this indicated that these isolates were of MANU1 lineage.

The current study reported on the isolation of MT strains from cattle from three different Indian states: Gujarat, Maharashtra and Telangana. The authors also reported on the genotyping of these strains, including those isolates from wild ruminants that had been described by Mukherjee *et al.* earlier (13). *Mycobacterium tuberculosis* was isolated from the lung tissues collected at necropsy of a gazelle (*G. bennettii*) and an antelope (*A. cervipara*) in 2010 and 2011, respectively. The animals were inhabitants of a nature park in the district of Gandhinagar, Gujarat. The nature park was home to, among other wild animals, seven gazelles and 150 antelopes that were housed in adjoining fenced enclosures, and free-ranging blue bulls (*Boselaphus tragocamelus*) ($n = 150$).

The nature park attendants lived in the villages surrounding the park located in Indroda, Gandhinagar, Gujarat, and also in villages surrounding Farm 9 in Gandhinagar, located 15 km from the nature park. (Farm 9 is the livestock farm in which the MT outbreaks in cattle referred to in this study occurred.) With the exception of core staff, the attendants left the park for their homes in the evening, and returned in the morning to resume their duties. The park had a record of persistent tuberculosis (TB), characterised by two episodes that resulted in mortality in eight deer in 2009–2010 followed by 29 deer in 2010–2011; necropsy of three of these wild ruminants revealed lesions suggestive of TB and impression smears from these samples tested positive using Acid Fast staining. The representative samples collected from the lesions at necropsy during both episodes, as reported by Mukherjee *et al.* (13), indicated that both outbreaks were due to MT infection since only *M. tuberculosis* was isolated, but not *M. bovis*. The main aims of this study were to attempt to:

a) isolate TB from cattle/buffaloes/wild ruminants sourced from farms/abattoirs/nature parks

b) characterise the isolates to the genotype level

c) explore the epidemiological significance of the identified genotypes in the animals examined

d) examine whether a correlation existed between the genotypes identified in the animals and those reported from human beings during the same period, from the same or neighbouring geographical locations.

In the present study, the significance of the results retrieved in pursuit of the abovementioned four aims is discussed.

Materials and methods

Animals and samples

Samples ($n = 841$) were collected from 733 adult female cattle, and 92 male cattle (31 adult, 61 young animals) and 16 buffaloes of one to 12 years of age from the three states (Telangana [$n = 352$], Maharashtra [$n = 31$] and Gujarat [$n = 458$]) from 2010 to 2015. Cattle below one year of age were considered young and those above one year of age as adult, while buffaloes less than two years of age were considered young and those above two years of age as adult. Out of the total of 841 samples, 405 were milk samples, 388 were nasal swabs, and the remaining 48 samples were of lymph nodes or liver lesions collected during post-mortem. The basis of sample collection and details associated with sample collection are described in Table I.

Specifically, a nasal swab (pooled from both nostrils) and a milk sample (pooled from all four quarters) was collected from each female animal that was in milk. For those female animals that were not in milk, only a pooled nasal swab was collected. On all but one occasion, only pooled nasal swabs were collected from male animals on farms; in the case of the exception, a pooled nasal swab and lymph node tissue were collected as samples. Lymph nodes were collected from

abattoirs as samples since, apart from the exception, no visible lung lesions were found during inspection after slaughter. (In the case of the exception, an abscess was found in the liver of an animal in a Mumbai abattoir.) This method of sampling was followed on 407 occasions, when it was only possible to collect a single sample from an animal in abattoirs and farms (Abattoir 1, $n = 16$; Farm 2, $n = 13$; Farm 3, $n = 16$; Farm 5, $n = 59$; Farm 6, $n = 62$; Abattoir 2, $n = 31$; Farm 7, $n = 90$; Farm 8, $n = 114$ and Farm 9, $n = 6$). On the remaining 434 occasions, it was possible to collect more than one sample from each animal (Table I). Table I also provides the status of samples with respect to: *a*) the presence or absence of a recorded clinical history of TB; and *b*) the completion status of a single intradermal tuberculin test (SITT). The tissues from necropsy and milk samples were collected in sterile 50 ml Falcon tubes, and nasal swabs were transferred to sterile Middlebrook 7H9 broth.

Interpretation of results of the single intradermal tuberculin test

The single intradermal tuberculin test (SITT) was conducted as described previously by the World Organisation for Animal Health (OIE) and Mukherjee *et al.* (1, 13) employing bovine tuberculin purified protein derivative (PPD) from Prionics (Schlieren, Switzerland). The SITT results were interpreted according to the criteria described in previous publications (1, 13). In brief, 72 h after inoculation with bovine tuberculin PPD, the skin-fold thickness of each animal was measured at the site of inoculation. An increase in the skin thickness equal to or greater than 4 mm, with or without inflammatory signs at 72 h, compared with the reading prior to inoculation, was interpreted as a positive test result.

Isolation of *Mycobacterium* from clinical samples

The milk samples were decontaminated and processed as previously described by Gao *et al.* (18). Briefly, 50 ml of milk samples were centrifuged at $3,100 \times g$ for 30 min to collect a pellet. The pellet and cream layer were decontaminated using hexadecylpyridinium chloride (HPC) at a final concentration of 0.75% for 5 h at 22°C. Next, samples

were centrifuged at $1,000 \times g$ for 15 min at room temperature, after which the supernatant was decanted (18). Nasal swabs and tissue samples collected during post-mortem were decontaminated and processed using a modified version of Petroff's method. In short, tissue samples and nasal samples were decontaminated through the addition of an equal volume of 4% sodium hydroxide, and were allowed to stand for 20 min. Next, they underwent two washings with 10 ml of sterile phosphate buffer solution (PBS) at pH 6.8 and finally the samples were centrifuged at $3,000 \times g$ for 15 min, and the supernatant decanted (19). Sediments obtained following decontamination of the milk samples, nasal swabs and tissues, as described above, were suspended in 2.0 ml of Middlebrook 7H9 broth. From each decontaminated suspension, 0.5 ml was inoculated into a BD BACTEC™ MGIT™ (Mycobacterial Growth Indicator Tube) 960/320 7 ml Tube containing 7 ml of modified Middlebrook 7H9 broth base already incorporated with BD BACTEC™ MGIT™ 960/320 Growth Supplement (GS) and 0.8 ml of the lyophilised antibiotic mixture – PANTA reconstituted with GS from BD BACTEC™ MGIT™ 960/320 Growth Supplement kit (Becton Dickinson [BD], Wokingham, United Kingdom) and the tubes were incubated in a BD BACTEC™ MGIT™ 960 system for 49 days at 37°C. In each case, from the remaining suspension, three tubes were inoculated: *a*) 0.1 ml was placed in a tube containing albumin dextrose catalase (OADC)-supplemented BD-DIFCO™ Middlebrook 7H10 agar (BD); *b*) 0.1 ml in a tube containing Lowenstein-Jensen (LJ) medium (HiMedia, Mumbai, India) supplemented with glycerol, and *c*) 0.1 ml in a tube containing LJ medium supplemented with sodium pyruvate; and these inoculated tubes containing three different media were incubated for eight weeks at 37°C in an ordinary air incubator.

Identification of *Mycobacterium* from culture

Heat-fixed smears prepared from MGIT cultures declared as positive by the BD BACTEC™ MGIT™ 960 system and typical growths on Middlebrook 7H10 and LJ media were screened for the presence of acid fast bacilli (AFB). The heat-fixed smears were stained for AFB using two commercial staining kits (TB Quick Stain Kit [BD, India,

Gurgaon, India], for the identification of AFB, and the TB Fluorescent Stain Kit M, using auramine-rhodamine [BD, India]). The smears were processed for staining according to the manufacturer's instructions.

Molecular identification of acid fast bacilli positive isolates

DNA extraction from MGIT liquid culture and colonies on Middlebrook 7H10 agar/LJ media was performed according to the protocol recommended for the GenoType® MTC kit (Hain Lifescience, Nehrem, Germany). To identify isolates as members of the MTC, the extracted DNA was amplified according to the protocol specified for the single-tube-nested polymerase chain reaction (PCR) kit (GeNei, Bangalore, India). In the single-tube two-step assays, the first positive amplification was determined by a 220-base pair (bp) PCR product amplified from the *IS6110* region, followed by amplification of a 123-bp-nested amplicon. For the identification and confirmation of strains, three commercial line-probe assay kits recommended by the World Health Organization (WHO) based on the polymorphism in the *gyrB* gene were employed (Genotype® MTC, Genotype® Mycobacterium CM and Genotype® Mycobacterium AS; Hain Lifescience). As well as identifying the presence of MTC, these kits can differentially identify *M. tuberculosis* from other members of the MTC. The Genotype® MTC kit can differentiate *M. tuberculosis*/*M. cannetti*, *M. africanum*, *M. microti*, *M. bovis*, *M. caprae* and Bacillus Calmette-Guérin (BCG) strains of the MTC from each other. The Genotype® Mycobacterium CM kit detects 'Common Mycobacteria' (CM) including *M. avium* complex (MAC), and 27 clinically relevant mycobacteria that are classified as non-tuberculous mycobacterium (NTM) or as mycobacterium other than tuberculosis (MOTT). The Genotype® Mycobacterium AS kit detects 'additional species' (AS) of *Mycobacteria* that includes another 19 clinically relevant MOTTs/NTMs, but not those that belong to MAC. In this study, the MTC kit was used as the primary test and the CM kit was used to double-check or reconfirm that the isolates really belonged only to the MTC group, whereas the AS kit was used to out rule the presence of NTMs/MOTTs in cultures.

Spoligotyping of *Mycobacterium* strains isolated from domestic and wild ruminants

Primers DRa (0.2 $\mu\text{mol}/\mu\text{l}$) and DRb (0.2 $\mu\text{mol}/\mu\text{l}$) were used for spoligotyping (20). The spacers between the direct repeats in the target region were amplified using two 18-nucleotide primers (primer 5'-CCAAGAGGGGACGGAAAC-3' and biotinylated primer 5'-GGTTTTGGGTCTGACGAC-3'). The PCR products were then hybridised to a membrane (Ocimum Biosolutions, Hyderabad, India). Hybridised DNA was detected using an enhanced chemiluminescence kit (Bio Basic, Israel), with exposure to X-ray film producing a pattern or profile reminiscent of a barcode. The hybridisation pattern was analysed in SPOTCLUST using the SpolDB3-based Model (17). SPOTCLUST 'clusters' spoligotype data for tuberculosis using mathematical models for genotyping, incorporating biological information on spoligotype evolution and epidemiology gained from epidemiological data (<http://tbinsight.cs.rpi.edu>).

Epidemiological analysis of *Mycobacterium tuberculosis* strains from ruminants with respect to prevailing human strains

Since a check on the *Mycobacterium tuberculosis* molecular markers database (SITVIT) for the origin and distribution of the characterised strains showed that they correlated with an Indian origin of strains, the ruminant MT spoligotypes were compared with human strains from India, using information available from the period of 2004 to 2017 (21, 22, 23, 24, 25, 26, 27, 28, 29) to create a unified information base for a region- and period-specific comparison. The results of the drug sensitivity test (DST) of the human MT strains of Indian origin belonging to various spoligotype lineages published previously (22, 27, 28, 29, 30) were analysed for information on the susceptibility or resistance of these strains to drugs.

Results

Isolation of *Mycobacterium* strains and their confirmation as *Mycobacterium tuberculosis*

Mycobacteria were isolated from seven of the 841 samples, comprising 4/405 milk, 2/388 nasal swabs and 1/48 post-mortem tissue samples from all the three Indian states examined (Gujarat, Maharashtra and Telangana) in MGITs, on Middlebrook 7H10 agar, and on LJ medium supplemented with glycerol, but not on LJ media supplemented with sodium pyruvate. Of the seven isolates, three were recovered from Farm 6 in Telangana, two from Farm 9 in Gujarat, one from Farm 4 in Telangana and the remaining isolate from an abattoir in Maharashtra (Tables I and II). Smears of the growth obtained in every culture system used showed the presence of slender acid-fast bacilli upon staining. Nested PCR of DNA from the isolates yielded both the 220- and 123-bp amplicons characteristic of MTC members. The hybridisation pattern generated by the Genotype® MTC kit identified all seven isolates as MT. The results of Genotype® Mycobacterium CM kit reconfirmed the isolates as members of the MTC and ruled out the presence of *M. avium* strains, while the Genotype® Mycobacterium AS kit confirmed the absence of MOTT strains (Tables I and II). The results of identification by culture, nested PCR and hybridisation pattern generated by the three commercial line-probe assay kits did not indicate the presence of any other member of the MTC (*M. bovis*, *M. bovis* BCG, *M. caprae*, *M. cannetti*, *M. africanum* or *M. microti*) except *M. tuberculosis*. Moreover, none of the battery of tests described above indicated the presence of members of *Mycobacteria* belonging to the MAC or MOTT groups.

Spoligotyping of *Mycobacterium tuberculosis* strains using SPOTCLUST

The spoligotyping analysis indicated the presence of MT strains belonging to the East African Indian (EAI) 5 family (Octal code – 47437777413771) from cattle and a gazelle (Octal code – 45437777413761) in Gujarat. The EAI3_IND family was identified in cattle and an antelope from Gujarat, and also from a liver lesion

from a male adult bovine in Maharashtra (Octal code – 47777777413071). Three distinct families, U (Octal code – 477776077411771), CAS1_DELHI (Octal codes – 703777740000771 and 702777340000571) as well as T1 (Octal code – 77771777760761), were identified from cattle in Telangana (Tables II and III).

Epidemiological analysis of *Mycobacterium tuberculosis* strains in ruminants and humans

The results of the study indicated the presence of the same spoligotypes in ruminants and humans living in the same and adjoining geographical region during the period under study and seemed to suggest the existence of an active ‘spillover’ mechanism of MT infection from human to domestic and wild ruminants. The evidence for this is as follows: between 2010 and 2012, human MT isolates were identified as belonging to the EAI5 family in Gujarat (21) and in Mumbai in the adjoining western state of Maharashtra (22, 23). Moreover, the same EAI5 family was identified from a sample from a lung lesion take from a gazelle (*G. bennettii*) in 2010 and from nasal swabs of cattle in 2011 in Gujarat. Similarly, the EAI3_IND family was reported from human MT isolates in Gujarat in 2010–2011 (21) from extra-pulmonary lesion in Mumbai, Maharashtra, in 2012 (23), and, in the present study, EAI3_IND was identified in an antelope (*A. cervipara*) and in cattle in Gujarat in 2011 and 2010, respectively.

In addition, the same spoligotype was identified in a strain isolated from a bovine liver in Mumbai, Maharashtra, in 2012. Furthermore, the U family (ST 1429 and ST 124) was reported in humans from Bhopal, Madhya Pradesh, during 2007–2011 (26), from Puducherry in 2015 (in extra-pulmonary lesions) (28), and from Vellore, Tamil Nadu, in 2017 (29). (The southern state of Tamil Nadu adjoins the state of Telangana in the north.) Kandhakumari *et al.*'s report, published in 2015 (28), was the first time that the U family had been identified from human strains in India. In this study, an MT isolate from cow's milk from Telangana was identified in 2010 as belonging

to the U family, probably indicating its prior presence in the southern region of India. *Mycobacterium tuberculosis* isolates from human pulmonary lesions, belonging to the CAS1_Delhi family, were reported in Tamil Nadu in 2010 (21) and Hyderabad, Telangana, in 2011 (27). The same family was identified from cow's milk samples from Hyderabad in 2010 and again in 2012. In addition, T1–T2 and T1 families were reported in human MT isolates from: the southern state of Tamil Nadu in 2010–2011, Bhopal in central India, Madhya Pradesh in 2007–2011, and Hyderabad, Telangana, in 2011 (21, 26, 27). In the present study, the T1 family was identified from nasal swabs of cattle in Hyderabad in 2011. Details of the above analysis are summarised in Table III. Drug sensitivity testing (DST) was not performed on isolates of animal origin in this study. Of the seven animals that were positive for MT, post-mortem examination could be conducted only on one animal; post-mortem examination revealed lesions in the liver.

Discussion

Natural infection of cattle, buffaloes and other ruminants with MB has been reported in India. *Mycobacterium bovis* has been isolated from lymph nodes (5), lung lesions (12), milk (14, 15, 16), and blood from cattle in the states of Himachal Pradesh (5, 12), Uttarakhand (14), Uttar Pradesh (15) and several other states in northern India (16), as mentioned in the reports published from the period beginning from 2005 to 2015. However, the concurrent infection of cattle with MB and MT has been also reported in India (12, 14, 15, 16). Against this background, infection with MT alone, in animals belonging to the Bovidae family is rarely reported in this country (11, 31, 32). Therefore, the authors were surprised to find only MT present in all seven isolates in this study, as well as in the wild ruminants previously reported by Mukherjee *et al.* in 2015 (13). These findings become more significant when one considers that MT was isolated from three species of ruminants (cattle, antelope and gazelle) from various geographical regions of India.

In this report, the authors used the highly sensitive and recommended commercial liquid culture system, BD BACTEC™ MGIT™ 960 system (33, 34), as well as reverse slot blot line probe *Mycobacterium* species identification assays (35) for the isolation and confirmation of seven MT isolates collected from different locations in India. The identification and confirmation of MT infection alone in the different types of samples (milk, nasal swab and liver), originating from diverse geographical locations in India had not previously been reported. The isolation from ruminants of *M. avium* and those strains grouped as MOTT has been described earlier (35, 36, 37). To examine this possibility, the authors used two commercial line probe assay kits to rule out the presence of *M. avium* and MOTT strains, and the existence of a concurrent infection from the source.

In 2012, Thakur *et al.* reported on the identification of an MT strain in cattle from the northern Indian state of Himachal Pradesh in 2006 using the spoligotyping approach (12); interpretation of the spoligotype patterns by the authors indicated that two of the MT isolates belonged to MANU1. In the present study, the authors report, for the first time, on the presence and the regional distribution of different MT spoligotypes in cattle, from 2010 to 2015. In addition, the authors also present spoligotyping data on MT isolates from captive wildlife ruminants, *G. bennettii* and *A. cervipara*. While regional evidence seems to indicate the persistence of spoligotype lineages of East African Indian origin – EAI5 and EAI3_IND family – in the western states of Gujarat, and EAI3_IND in Maharashtra. A different set of lineages represented as CAS1_Delhi, U and T1 spoligotypes were identified in the southern state of Telangana in the period under study. The authors also report the simultaneous existence of two different MT cattle isolates with spoligotype lineages EAI3_IND and EAI5 in the same location in Gujarat (Farm 9 – Table I) responsible for tuberculosis outbreaks in 2010 and 2011, respectively. Similarly, the authors noticed the existence of two different spoligotypes in two different ruminant species identified as EAI5 in *G. bennettii* and EAI3_IND in *A. cervipara* from the same Nature Park in Gujarat (Laboratory identification No. MCR1 and MCR2 – Tables II and III) in 2010 and 2011, respectively. Since five

MT spoligotypes from seven ruminants in India were identified in this study, it is possible that more MT strains exist in ruminants in the subcontinent. A similar observation has previously been made based on the existence of three *M. bovis* spoligotypes in cattle in Ethiopia in just 11 cattle (7).

Although there is an abundance of information available related to the genotypes of human strains of MT prevalent in India (21, 22, 23, 24, 25, 26, 27, 28, 29, 31), similar reports on ruminants from this country are scarce. Only two previous reports (12, 13) have identified the existence of different genotypes of the MT strain in ruminants, so overall, including the current study, six MT lineages (MANU1, EAI3_IND, EAI5, CAS1_Delhi, U and T) have been detected from ruminants to date from a total of just three studies. Moreover, recently, in January 2017, MT isolates from cattle of the MANU1 lineage were reported in Tamil Nadu (32). This preliminary information derived from the limited number of studies in ruminants does not reveal the complete picture regarding the number of human strains of MT that exist in India's entire ruminant population. It does, however, indicate that in order to reveal the actual situation, a large number of further studies would be necessary. This supposition is strengthened by previous reports of the existence of different MT genotypes in cattle herds (7, 34) and the fact that in high prevalence settings or of high TB burden, humans and animals living in close contact for a prolonged period are quite likely to acquire *M. tuberculosis* infection (7, 16, 34, 38, 39, 40) due to their genotypes being the same or closely related as result of spill-over or spill-back mechanisms (38). It is probably through this mechanism that ruminants/cattle acquire infection as there are several MT strains of human origin worldwide, whereas MT variants do not occur in cattle.

Mycobacterium tuberculosis infection in animals can spread from one geographical area to another, due to the unrestricted movement of infected animals or contact with infected humans. The chronology of MT outbreaks in different ruminant species recorded in the area where Farm 9 and the nature park are located (13) probably indicates that the same spoligotype lineage EAI3_IND that infected cattle in 2010,

infected *A. cervipara* in the nature park in 2011. Similar MT spoligotypes were observed when the EAI5 identified in *G. bennettii* in 2010 was recovered from cattle in Farm 9 in 2011. Large-scale unrestricted road migration of farmers/transients along with their sick, unproductive, infected and stressed animals, due to drought and shortage of fodder, is common from Gujarat to the adjacent state of Maharashtra. In this manner, infected animals gain access to a new location and new hosts. Such unproductive animals ultimately end up in the abattoirs of Mumbai. It seems, perhaps, that this could be one of the probable mechanisms through which MT infection in cattle, reported from Gandhinagar, Gujarat in 2010 due to EAI3_IND lineage, spread to Mumbai, Maharashtra, in 2012. In contrast, the authors observed two different MT strains, U and CAS1_Delhi, in the same farm (Farm 6) in Telangana also in 2010. Moreover, yet another spoligotype variant, T1, was reported in another farm (Farm 4) in Telangana in 2011. The reason for the predominant presence of CAS spoligotype in humans in Hyderabad in recent times, a strain mostly reported from northern India, could be due to the increasing influx and convergence of non-resident Indians (NRI) working in the Middle East and the migrant population from northern India which now reside in Hyderabad, the commercial capital of Telangana (38). This study's observation of the existence of CAS1_Delhi spoligotype in cattle in 2010 could be due to the spill-over effect from humans.

In the study of tuberculosis, India stands out as being home to *M. tuberculosis* lineage 1 (Indo-Oceanic or EAI lineage) and lineage 3 (Central Asian or CAS lineage) primarily, which occur at substantially lower frequencies outside the Indian subcontinent. In contrast, lineages 2 (East Asian or Beijing) and 4 (Euro–American) are the most common lineages in Europe, Africa and many other parts of the world. Even within India, the prevalence of lineages varies. Lineage 3 predominates among patients from north and north-west India, and lineage 1 is commonly found among Southern Indian patients and at low frequency among patients in other parts of the country. In contrast, lineage 2 has been reported at similar prevalences throughout India, though it does predominate in some north-eastern Indian states (41, 42, 43).

Humans are the maintenance hosts for MT; however, cattle and wild ruminants have been identified as spill-over hosts (34, 38, 39). It has been observed that spill-over hosts are commonly infected when the level of challenge is relatively high (3). A high TB burden in humans infected with MT lineages, such as EAI3_IND, EAI5 and CAS1_Delhi prevailing in India may therefore be the reason for such a spillover to ruminants (44). At least, on four occasions, the isolation of MT from nasal swabs from cattle in this study (Table II) and from lung samples from wild ruminants in the study by Mukherjee *et al.* (13) seems to indicate the transmission of MT infection from humans to ruminants through an aerogenous route. This fact is in agreement with earlier observations (45, 46). Moreover, it was also evident in the current study, that MT was isolated from milk on at least four occasions, and from liver on one occasion (Table II), probably indicating acquired extra-pulmonary MT infection by humans. Transmission of MT infection from humans to 'spillover' hosts, such as cattle, from extra-pulmonary sources has been suggested previously (13, 16). Although the precise mechanism cannot be fully explained, the lineages of human MT strains isolated from extra-pulmonary tissue sources (19, 43) and the identification of EAI3_IND from extra-pulmonary lesions from cattle in this study strengthens this contention. Similarly, in recent times two separate studies from Ethiopia have indicated that farmers and their cattle were infected with MT (7, 40). Both these studies revealed that cattle, in addition to being infected with *M. bovis* and NTM, were also infected with MT strains that belonged three families – CAS, T3-ETH and T1 (7, 40, 47, 48).

The WHO *Global Tuberculosis Report 2015* indicated that India has the largest number of TB cases in the world with 23% (49). Compared to other regions in the world, MT infection reported in cattle in India over the last 12 years, with respect to isolation from samples is quite high, varying from 15% to 84% (32, 50). Since SITT is not able to discriminate MT from MB infection in cattle, and facilities for the culture of pathogenic *Mycobacterium* species are mostly not available in the rural settings in India, over all, the above facts give rise to a crucial diagnostic disadvantage that impacts the control of bovine tuberculosis in India.

The report on the sharing of identical MT strains of the same lineage in humans and cattle has been emphasised as an important epidemiological event that should be considered when trying to minimise the risk of the transmission of MT from humans to cattle and from cattle to humans by spillover and spillback mechanisms (34, 38). Earlier reports of the isolation of MT from bovine milk (11, 16, 50) and from extra-pulmonary lesions in cattle (50) and humans (28, 44), and the practices of drinking unpasteurised and soured milk, and of raising herds of cattle on open pastures in rural India, has brought the authors closer to the realisation that MT infection in cattle could pose a new challenge in the control of TB in humans in this country. Taken together, the above observations can impact the control of bovine and human TBs.

The relevance of the observed association between virulence intensity in cattle infected with *M. bovis* (51) and *M. tuberculosis* in humans (52) with certain MT genotypes has been highlighted. Since several reports, as discussed above (7, 16, 34, 38, 39, 50), have indicated that MT strains of the same family are shared among infected ruminants and humans, it would seem that further investigation is necessary to determine whether such MT genotypes are present in the ruminants in India. In the present study, it was possible to conduct a post-mortem on one animal in a Mumbai abattoir which showed a lesion in its liver; however, the authors were legally restricted from performing the same on six other MT bacterial culture positive live cattle, owned by private farms (Farms 4, 6 and 9), and therefore, were unable to associate the extent and intensity of lesions with culture positivity.

The authors observations on the distribution of MT strains of ruminant origin reported from the regions of Gandhinagar in Gujarat, Mumbai in Maharashtra and Hyderabad in Telangana (formerly in Andhra Pradesh) during 2010 to 2012, when compared to reports on the molecular epidemiology of human strains from 2004 to 2015 (21, 28), probably reflects the parallel existence of MT strains of same family lineage, in the human maintenance host and the spillover ruminant host. The simultaneous existence of the same MT families in cattle and in their herdsmen has been recently observed in farms in the

southern state of Tamil Nadu (P. Kannan, personal communication). The above observations when combined seem to suggest that a spill over situation of MT strains exists from humans to ruminants on the Indian sub-continent. Drug sensitivity tests of human MT strains recovered from ruminants were not undertaken in this study. However, the availability of reports on such tests may add a new dimension to the understanding of the epidemiology of MT infection in ruminants; the DST reports on human strains from India that are available, indicate a variance in the resistance or sensitivity to antibiotics (22, 27, 28, 29, 30). While pan-sensitivity of CAS1_DELHI (ST 26) and EAI5 (ST 236) was observed in one report (22), for example, in another the majority of human MT strains from India of EAI5, EAI3_IND and CAS families were resistant to multiple drugs (27).

Conclusions

The above observations certainly complicate the method of controlling a zoonotic disease, such as TB, in India, since MT strains seem to exist and cross-infect hosts without a species barrier at the same time point. It will be worth investigating the spill-back behaviour of these ruminant MT strains to humans, as reported in this study. Furthermore, a more intensive molecular epidemiological investigation on the presence of the MT strains prevalent in domestic and wild ruminants on the Indian sub-continent, other than those reported herein, is warranted in order to enhance clarity on the magnitude of the spillover effect on ruminants that prevails in India.

Acknowledgements

The authors are grateful to the management of the National Dairy Development Board in Anand for providing the facilities to carry out this work, which was conducted at the Research and Development (R&D) Laboratory, National Dairy Development Board (NDDB), Hyderabad. Mr Vijay Bahekar expresses his gratitude to the NDDB, for providing him with the opportunity to work on the isolation and

characterisation of *Mycobacterium* isolates as part of his PhD thesis at the R&D facilities of NDDDB at Hyderabad.

References

1. World Organisation for Animal Health (OIE) (2015). – Chapter 2.4.6. Bovine tuberculosis. *In* Manual of Diagnostic Tests and Vaccines for Terrestrial Animals. OIE, Paris, France, 17 pp. Available at:

www.oie.int/fileadmin/Home/eng/Health_standards/tahm/2.04.06_BOVINE_TB.pdf (accessed on 3 November 2017).

2. Cosivi O., Grange J.M., Daborn C.J., Raviglione M.C., Fujikura T., Cousins D., Robinson R.A., Huchzermeyer H.F.A.K., de Kantor I. & Meslin F.-X. (2008). – Zoonotic tuberculosis due to *Mycobacterium bovis* in developing countries. *Emerg. Infect. Dis.*, **4** (1), 59–70. doi:10.3201/eid0401.980108.

3. Cousins D.V. (2001). – *Mycobacterium bovis* infection and control in domestic livestock. *In* Mycobacterial infections in domestic and wild animals (E.J.B. Manning & M.T. Collins, eds). *Rev. Sci. Tech. Off. Int. Epiz.*, **20** (1), 71–85. doi:10.20506/rst.20.1.1263.

4. Firdessa R., Tschopp R. [...] & Berg S. (2012). – High prevalence of bovine tuberculosis in dairy cattle in central Ethiopia: implications for the dairy industry and public health. *PloS ONE*, **7** (12), e52851. doi:10.1371/journal.pone.0052851.

5. Thakur A., Sharma M., Katoch V.C., Dhar P. & Katoch R.C. (2010). – A study on the prevalence of bovine tuberculosis in farmed dairy cattle in Himachal Pradesh. *Vet. World*, **3** (9), 408–413. doi:10.5455/vetworld.2010.408-413l.

6. Jia K., Yu M., Zhang G.-H., Zhang J., Lin Z.-X., Luo C.-B., Yu H.-Q. & Li S.-J. (2012). – Detection and identification of *Mycobacterium tuberculosis* and *Mycobacterium bovis* from clinical species using DNA microarrays. *J. Vet. Diagn. Invest.*, **24** (1), 156–160. doi:10.1177/1040638711417141.

7. Tsegaye W., Aseffa A., Mache A., Mengistu Y., Stefan B. & Ameni G. (2010). – Conventional and molecular epidemiology of bovine tuberculosis in dairy farms in Addis Ababa city, the capital of Ethiopia. *Intern. J. Appl. Res. Vet. Med.*, **8** (2), 143–151. Available at: www.jarvm.com/articles/Vol8Iss2/Vol8%20Iss2Tsegaye.pdf (accessed on 3 November 2017).

8. Tuzcu N., Kayar B., Uysal E.B., Gülcü Y., Marzi M. & Köksal F. (2015). – Spoligotyping of *M. tuberculosis* strains from cattle in Turkey. *Kafkas Univ. Vet. Fak. Derg.*, **21** (4), 507–511. doi:10.9775/kvfd.2014.12769

9. Cadmus S., Palmer S., Okker M., Dale J., Gover K., Smith N., Jahans K., Hewinson R.G. & Gordon S.V. (2006). – Molecular analysis of human and bovine tubercle bacilli from a local setting in Nigeria. *J. Clin. Microbiol.*, **44** (1), 29–34. doi:10.1128/JCM.44.1.29-34.2006.

10. Romero B., Rodríguez S. [...] & de Juan L. (2011). – Humans as source of *Mycobacterium tuberculosis* infection in cattle, Spain. *Emerg. Infect. Dis.*, **17** (12), 2393–2395. doi:10.3201/eid1712.101476.

11. Bhanu Rekha V., Gunaseelan L., Pawar G., Nassiri R. & Bharathy S. (2015). – Molecular detection of *Mycobacterium tuberculosis* from bovine milk samples. *J. Adv. Vet. Anim. Res.*, **2** (1), 80–83. doi:10.5455/javar.2015.b44.

12. Thakur A., Sharma M., Katoch V.C., Dhar P. & Katoch R.C. (2012). – Detection of *Mycobacterium bovis* and *Mycobacterium tuberculosis* from cattle: possible public health relevance. *Indian J. Microbiol.*, **52** (2), 289–291. doi:10.1007/s12088-011-0200-8.

13. Mukherjee F., Bahekar V.S., Prasad A., Rana S.K., Kanani A., Sharma G.K. & Srinivasan V.A. (2015). – Isolation of *Mycobacterium tuberculosis* from *Antelope cervicapra* and *Gazelle bennettii* in India and confirmation by molecular tests. *Eur. J. Wildl. Res.*, **61** (5), 783–787. doi:10.1007/s10344-015-0938-0.

14. Kumar M., Kumar A., Sharma N. & Singh R.K. (2015). – Molecular characterization of *M. tuberculosis* and *M. bovis*-clinical application for the disease management. *J. Sci.*, **5** (5), 300–303. Available at: www.journalofscience.net/doi/MjI1a2FsYWkxNDc4NTIzNjk= (accessed on 22 November 2017).

15. Mishra A., Singhal A., Chauhan D.S., Katoch V.M., Srivastava K., Thakral S.S., Bharadwaj S.S., Sreenivas V. & Prasad H.K. (2005). – Direct detection and identification of *Mycobacterium tuberculosis* and *Mycobacterium bovis* in bovine samples by a novel nested PCR assay: correlation with conventional techniques. *J. Clin. Microbiol.*, **43** (11), 5670–5678. doi:10.1128/JCM.43.11.5670-5678.2005.

16. Srivastava K., Chauhan D.S., Gupta P., Singh H.B., Sharma V.D., Yadav V.S., Sreekumaran, Thakral S.S., Dharamdheeran J.S., Nigam P., Prasad H.K. & Katoch V.M. (2008). – Isolation of *Mycobacterium bovis* and *M. tuberculosis* from cattle of some farms in north India – possible relevance in human health. *Indian J. Med. Res.*, **128** (1), 26–31. Available at: www.semanticscholar.org/paper/Isolation-of-Mycobacterium-bovis-%26-M.-tuberculosis-Srivastava-Chauhan/0f910fb5d86dc14c3a814727e5399840bc026974 (accessed on 11 October 2018).

17. Vitol I., Driscoll J., Kreiswirth B., Kurepina N. & Bennett K.P. (2006). – Identifying *Mycobacterium tuberculosis* complex strain families using spoligotypes. *Infect. Genet. Evol.*, **6** (6), 491–504. doi:10.1016/j.meegid.2006.03.003.

18. Gao A., Odumeru J., Raymond M. & Mutharia L. (2005). – Development of improved method for isolation of *Mycobacterium avium* subsp. *paratuberculosis* from bulk tank milk: effect of age of milk, centrifugation, and decontamination. *Can. J. Vet. Res.*, **69** (2), 81–87. Available at: www.ncbi.nlm.nih.gov/pmc/articles/PMC1142174/ (accessed on 3 November 2017).

19. Kent P.T. & Kubica G.P.W. (1985). – Public health mycobacteriology. A guide for the level III Laboratory. Centers for Disease Control and Prevention (CDC), Atlanta, Georgia, United States of America, 21–44.

20. Kamerbeek J., Schouls L., Kolk A., van Agterveld M., van Soolingen D., Kuijper S., Bunschoten A., Molhuizen H., Shaw R., Goyal M. & van Embden J. (1997). – Simultaneous detection and strain differentiation of *Mycobacterium tuberculosis* for diagnosis and epidemiology. *J. Clin. Microbiol.*, **35** (4), 907–914. Available at: www.ncbi.nlm.nih.gov/pmc/articles/PMC229700/ (accessed on 3 November 2017).

21. National Institute for Research in Tuberculosis (NIRT) (2011). – Profiling of molecular heterogeneity and identification of region-specific gene segments in the field strains of *M. tuberculosis*. In National Institute for Research in Tuberculosis: research activities. April 2010–March 2011 [Annual Report 2010–2011]. NIRT, Chennai, India, 69–76. Available at: <http://nirt.res.in/html/annual%20reports.html> (accessed on 11 October 2018).

22. Chatterjee A., D'Souza D., Vira T., Bamne A., Ambe G.T., Nicol M.P., Wilkinson R.J. & Mistry N. (2010). – Strains of *Mycobacterium tuberculosis* from western Maharashtra, India, exhibit a high degree of diversity and strain-specific associations with drug resistance, cavitary disease, and treatment failure. *J. Clin. Microbiol.*, **48** (10), 3593–3599. doi:10.1128/JCM.00430-10.

23. Vadwai V., Shelly A., Supply P. & Rodrigues C. (2012). – Evaluation of 24-locus MIRU-VNTR in extrapulmonary specimens: study from a tertiary centre in Mumbai. *Tuberculosis (Edinb.)*, **92** (3), 264–272. doi:10.1016/j.tube.2012.01.002.

24. National Institute for Research in Tuberculosis (NIRT) (2009). – Epidemiological and operational research: completed studies. Spoligotyping *M. tuberculosis* isolates from Tiruvallur district. In Tuberculosis Research Centre: research activities. April

2008–March 2009 [NIRT Annual Report 2008–2009]. NIRT, Chennai, India, 39–41. Available at: www.nirt.res.in/pdf/ar/trc-ar-2008-2009.pdf (accessed on 6 November 2017).

25. Indian Council of Medical Research (ICMR) (2014–2015). – Communicable diseases. Genetic diversity and molecular characterisation of drug resistance in *Mycobacterium tuberculosis* isolates from south India by whole genome sequencing. In Annual report 2014–15, ICMR, New Delhi, India, 2 and Fig 1. Available at: <https://icmr.nic.in/icmr-annual-reports> (accessed on 26 October 2018).

26. Desikan P., Chauhan D.S., Sharma P., Panwalkar N., Chourey M., Patidar M.L., Yadav P., Chandrasekaran V. & Ohri B.S. (2016). – Genetic diversity of *Mycobacterium tuberculosis* isolates from central India. *Indian J. Med. Res.*, **143** (4), 481–486. doi:10.4103/0971-5916.184287.

27. Thomas S.K., Iravatham C.C., Moni B.H., Kumar A., Archana B.V., Majid M., Priyadarshini Y., Rani P.S., Valluri V., Hasnain S.E. & Ahmed N. (2011). – Modern and ancestral genotypes of *Mycobacterium tuberculosis* from Andhra Pradesh, India. *PLoS ONE*, **6** (11), e27584. doi:10.1371/journal.pone.0027584.

28. Kandhakumari G., Stephen S., Sivakumar S. & Narayanan S. (2015). – Spoligotype patterns of *Mycobacterium tuberculosis* isolated from extra pulmonary tuberculosis patients in Puducherry, India. *Indian J. Med. Microbiol.*, **33** (2), 267–270. doi:10.4103/0255-0857.154871.

29. Suzana S., Shanmugam S., Uma Devi K.R., Swarna Latha P.N. & Michael J.S. (2017). – Spoligotyping of *Mycobacterium tuberculosis* isolates at a tertiary care hospital in India. *Trop. Med. Int. Hlth*, **22** (6), 703–707. doi:10.1111/tmi.12875.

30. Manson A.L., Abeel T., Galagan J.E., Sundaramurthi J.C., Salazar A., Gehrman T., Shanmugam S.K., Palaniyandi K., Narayanan S., Swaminathan S. & Earl A.M. (2017). – *Mycobacterium tuberculosis* whole genome sequences from southern India suggest

novel resistance mechanisms and the need for region-specific diagnostics. *Clin. Infect. Dis.*, **64** (11), 1494–1501. doi:10.1093/cid/cix169.

31. Mukhopadhyay S.K. (2007). – Studies on prevalence and aetiopathological aspect of common diseases in spotted deer in West Bengal. *Indian J. Vet. Pathol.*, **31** (1), 35–39.

32. Sweetline Anne N., Ronald B.S.M., Senthil Kumar T.M.A., Kannan P. & Thangavelu A. (2017). – Molecular identification of *Mycobacterium tuberculosis* in cattle. *Vet. Microbiol.*, **198**, 81–87. doi:10.1016/j.vetmic.2016.12.013.

33. Robbe-Austerman S., Bravo D.M. & Harris B. (2013). – Comparison of the MGIT 960, BACTEC 460 TB and solid media for isolation of *Mycobacterium bovis* in United States veterinary specimens. *BMC Vet. Res.*, **9**, 74. doi:10.1186/1746-6148-9-74.

34. Romero B., Rodríguez S. [...] & de Juan L. (2011). – Humans as source of *Mycobacterium tuberculosis* infection in cattle, Spain [letter]. *Emerg. Infect. Dis.*, **17** (12), 2393–2395. doi:10.3201/eid1712.101476.

35. Gortazar C., Torres M.J., Acevedo P., Aznar J., Negro J.J., de la Fuente J. & Vicente J. (2011). – Fine-tuning the space, time, and host distribution of mycobacteria in wildlife. *BMC Microbiol.*, **11**, 27. doi:10.1186/1471-2180-11-27.

36. Romero B., Aranaz A., Sandoval Á., Álvarez J., de Juan L., Bezos J., Sánchez C., Galka M., Fernández P., Mateos A. & Domínguez L. (2008). – Persistence and molecular evolution of *Mycobacterium bovis* population from cattle and wildlife in Doñana National Park revealed by genotype variation. *Vet. Microbiol.*, **132** (1–2), 87–95. doi:10.1016/j.vetmic.2008.04.032.

37. Thorel M.-F., Huchzermeyer H.F. & Michel A.L. (2001). – *Mycobacterium avium* and *Mycobacterium intracellulare* infection in mammals. In *Mycobacterial infections in domestic and wild animals*

(E.J.B. Manning & M.T. Collins, eds). *Rev. Sci. Tech. Off. Int. Epiz.*, **20** (1), 204–218. doi:10.20506/rst.20.1.1272.

38. Malama S., Muma J., Munyeme M., Mbulo G., Muwonge A., Shamputa I.C., Djønne B., Godfroid J. & Johansen T.B. (2014). – Isolation and molecular characterization of *Mycobacterium tuberculosis* from humans and cattle in Namwala district, Zambia. *EcoHealth*, **11** (4), 564–570. doi:10.1007/s10393-014-0940-0.

39. Michel A.L., Hlokwe T.M., Espie I.W., van Zijll Langhout M., Koepel K. & Lane E. (2013). – *Mycobacterium tuberculosis* at the human/wildlife interface in a high TB burden country. *Transbound. Emerg. Dis.*, **60** (S1), S46–S52. doi:10.1111/tbed.12099.

40. Ameni G., Tadesse K., Hailu E., Deresse Y., Medhin G., Aseffa A., Hewinson G., Vordermeier M. & Berg S. (2013). – Transmission of *Mycobacterium tuberculosis* between farmers and cattle in Central Ethiopia. *PLoS ONE*, **8** (10), e76891. doi:10.1371/journal.pone.0076891.

41. Narayanan S., Gagneux S., Hari L., Tsolaki A.G., Rajasekhar S., Narayanan P.R., Small P.M., Holmes S. & DeRiemer K. (2008). – Genomic interrogation of ancestral *Mycobacterium tuberculosis* from south India. *Infect. Genet. Evol.*, **8** (4), 474–483. doi:10.1016/j.meegid.2007.09.007.

42. Shanmugam S., Selvakumar N. & Narayanan S. (2011). – Drug resistance among different genotypes of *Mycobacterium tuberculosis* isolated from patients from Tiruvallur, South India. *Infect. Genet. Evol.*, **11** (5), 980–986. doi:10.1016/j.meegid.2011.03.011.

43. Singh J., Sankar M.M., Kumar P., Couvin D., Rastogi N. & Singh S. (2015). – Genetic diversity and drug susceptibility profile of *Mycobacterium tuberculosis* isolated from different regions of India. *J. Infect.*, **71** (2), 207–219. doi:10.1016/j.jinf.2015.04.028.

44. Sankar M.M., Singh J., Diana S.C.A. & Singh S. (2013). – Molecular characterization of *Mycobacterium tuberculosis* isolates

from North Indian patients with extrapulmonary tuberculosis. *Tuberculosis (Edinb.)*, **93** (1), 75–83. doi:10.1016/j.tube.2012.10.005.

45. Müller B., Dürr S., Alonso S., Hattendorf J., Laise C.J.M., Parsons S.D.C., van Helden P.D. & Zinsstag J. (2013). – Zoonotic *Mycobacterium bovis* – induced Tuberculosis in humans. *Emer. Infect. Dis.*, **19** (6), 899–908. doi:10.3201/eid1906.120543.

46. Dürr S., Müller B., Alonso S., Hattendorf J., Laise C.J.M., van Helden P.D. & Zinsstag J. (2013). – Differences in primary sites of infection between zoonotic and human tuberculosis: results from a worldwide systematic review. *PLoS Negl. Trop. Dis.*, **7** (8), e2399. doi:10.1371/journal.pntd.0002399.

47. Debebe T., Admassu A., Mamo G. & Ameni G. (2014). – Molecular characterization of *Mycobacterium tuberculosis* isolated from pulmonary tuberculosis patients in Felege Hiwot Referral Hospital, northwest Ethiopia. *J. Microbiol. Immunol. Infect.*, **47** (4), 333–338. doi:10.1016/j.jmii.2013.03.012.

48. Belay M., Ameni G., Bjune G., Couvin D., Rastogi N. & Abebe F. (2014). – Strain diversity of *Mycobacterium tuberculosis* isolates from pulmonary tuberculosis patients in Afar Pastoral region of Ethiopia. *BioMed. Res. Int.*, **2014**, Article ID 238532. doi:10.1155/2014/238532.

49. World Health Organization (WHO) (2015). – WHO/HTM/TB/2015.22. Global tuberculosis report 2015, 20th Ed., WHO, Geneva, Switzerland, 204 pp. Available at: http://apps.who.int/iris/bitstream/handle/10665/191102/9789241565059_eng.pdf?sequence=1 (accessed on 12 October 2018).

50. Prasad H.K., Singhal A., Mishra A., Shah N.P., Katoch V.M., Thakral S.S., Singh D.V., Chumber S., Bal S., Aggarwal S., Padma M.V., Kumar S., Singh M.K. & Acharya S.K. (2005). – Bovine tuberculosis in India: potential basis for zoonosis. *Tuberculosis (Edinb.)*, **85** (5–6), 421–428. doi:10.1016/j.tube.2005.08.005.

51. Garbaccio S., Macias A. [...] & Cataldi A. (2014). – Association between spoligotype-VNTR types and virulence of *Mycobacterium bovis* in cattle. *Virulence*, **5** (2), 297–302. doi:10.4161/viru.27193.

52. Caminero J.A., Pena M.J. [...] & Martin C. (2001). – Epidemiological evidence of the spread of a *Mycobacterium tuberculosis* strain of the Beijing genotype on Gran Canaria Island. *Am. J. Respir. Crit. Care Med.*, **164** (7), 1165–1170. doi:10.1164/ajrccm.164.7.2101031.

Table I
Details of the samples collected and processed for isolation from 2010 to 2015

Origin of sample (source, location, state)	Single intradermal tuberculin test (SITT) result ^(a)	Clinical history of tuberculosis (TB)	Type of sample ^(b)			No. of isolates (year of sampling and isolation)
			Milk	Nasal swab	Tissue	
Abattoir 1, Hyderabad, Telangana	Not done	Not available	0	0	16	0
Farm 1, Hyderabad, Telangana	0 (64)	No history of TB	64	64	0	0
Farm 2, Hyderabad, Telangana	Not done	No history of TB	13	0	0	0
Farm 3, Hyderabad, Telangana	Not done	No history of TB	4	20	0	0
Farm 4, Hyderabad, Telangana	Not done	History of TB	(1) 11	11	0	1 (2011)
Farm 5, Hyderabad, Telangana	Not done	No history of TB	0	60	1	0
Farm 6, Hyderabad, Telangana	18 (29)	History of TB	(3) 75	13	0	3 (2010, 2010, 2012)
Abattoir 2, Mumbai, Maharashtra	Not done	Not available	0	0	(1) 31	1 (2012)
Farm 7, Anand, Gujarat	0 (150)	No history of TB	60	150	0	0
Farm 8, Bidaj, Gujarat	0 (144)	No history of TB	144	30	0	0
Farm 9, Gandhinagar, Gujarat	Not done	History of TB	(1) 34	(1) 40	0	2 (2010 and 2011)
Total no. of samples: 841			405	388	48	7

a) Some or all animals tested in the farms were positive by the single intradermal tuberculin test (SITT) using bovine tuberculin purified protein derivative (PPD) from Prionics, Switzerland; positive and negative test results were interpreted according to the standard interpretation criteria (1, 13); where figures are included, the number of positive animals is followed by the total number of tested animals in parenthesis

b) The number of positive isolates from different types of samples, where available, is contained in parenthesis; 'tissues' denotes lymph nodes or liver lesions

Table II
Details of *Mycobacterium tuberculosis* field isolates from domestic and wildlife ruminants and spoligotyping results

Sample ID no.	Species and age	Gender	Sample	Remarks on isolation	Geographical location of isolates	Octal code	SpoIDB3 based lineage	Binary spoligotype patterns
MCR1	Gazelle, one year	Female	Lung	Previous study (13)	Gandhinagar, Gujarat	45437777413761	EAI5	
MCR2	Antelope, one year	Female	Lung	Previous study (13)	Gandhinagar, Gujarat	47777777413071	EAI3_IND	
MCR3	Cattle, six years	Female	Milk	This study	Farm 9, Gandhinagar, Gujarat	47777777413071	EAI3_IND	
MCR4	Cattle, eight years	Female	Milk	This study	Farm 6, Hyderabad, Telangana	47776077411771	U	
MCR5	Cattle, nine years	Female	Milk	This study	Farm 6, Hyderabad, Telangana,	703777740000771	CAS1-Delhi	
MCR6	Cattle, seven years	Female	Nasal swab	This study	Farm 9, Gandhinagar, Gujarat	47437777413771	EAI5	
MCR7	Cattle, ten years	Male	Liver	This study	Abattoir 2, Mumbai, Maharashtra	47777777413071	EAI3_IND	
MCR8	Cattle, seven years	Female	Milk	This study	Farm 6, Hyderabad, Telangana,	702777340000571	CAS1-Delhi	
MCR9	Cattle, eight years	Female	Nasal swab	This study	Farm 4, Hyderabad, Telangana	77771777760761	T1	

Open squares: lack of hybridisation; represented from 1 to 43

Solid squares: hybridisation with designated spacer probe

CAS: Central and Middle Eastern Asia

EAI: East African Indian

T1: Tuscany

U: unidentified

The *Mycobacterium tuberculosis* (MT) isolates in wild female ruminant species were reported during a previous study (13) and the spoligotyping results of these isolates were reported during this study. The MT isolates from cattle were identified and spoligotyped in this study. All isolates were recovered from cultures in the BACTEC™ MGIT™ 960 system (Beckton Dickinson, Wokingham, United Kingdom), BD-DIFCO™ Middlebrook 7H10 agar supplemented with oleic albumin dextrose catalase (OADC) and glycerol, and Lowenstein Jensen medium supplemented with glycerol but not sodium pyruvate, and were identified as members of the *Mycobacterium tuberculosis* complex (MTC) group by IS6110 commercial nested polymerase chain reaction (PCR) (GeNei, Bangalore, India). All isolates were confirmed as MT by a specific pattern of reverse slot blot hybridisation of the line probe assay using Genotype® MTC kit (Hain Lifescience, Nehrem, Germany). The spoligotype patterns were identified among MT isolates. Clade designations were made according to SpolDB3. Genotyping of all these isolates were performed in this study

Table III

Details of *Mycobacterium tuberculosis* field isolates from domestic and wildlife ruminants and spoligotyping results as compared to regional presence of human strains

Laboratory identification no.	Species ^{(a) (b)}	Tissue	Geographical location of isolates	Year of isolation	Clade ^(c)	Regional presence of families of <i>Mycobacterium tuberculosis</i> human isolates ^(d)			
						Spoligotype/clade	Year of isolation/reporting	Location, region / tissue	References
MCR1	Gazelle	Lung	Gandhinagar, Gujarat	2010	EAI5	EAI5	2010, 2011; 2012	Gujarat; Maharashtra, Mumbai; Maharashtra, Mumbai	(21, 22, 23)
MCR2	Antelope	Lung	Gandhinagar, Gujarat	2011	EAI3_IND	EAI3_IND	2010, 2011	Gujarat	(21)
MCR3	Cattle	Milk	Gandhinagar, Gujarat	2010	EAI3_IND	EAI3_IND	2010, 2011	Gujarat	(21)
MCR4	Cattle	Milk	Hyderabad, Telangana	2010	U	U	2007–2011, 2015	Bhopal, Madhya Pradesh; Tamil Nadu, Puducherry; Tamil Nadu, Vellore	(26, 28, 29)
MCR5	Cattle	Milk	Hyderabad, Telangana	2010	CAS1-Delhi	CAS1_Delhi	2011; 2008–2009; 2014–2015; 2010,	Tamil Nadu, Tamil Nadu, Tiruvallur; Southern India; Telangana (Andhra Pradesh) / pulmonary;	(21, 24, 25, 27)
MCR6	Cattle	Nasal swab	Gandhinagar, Gujarat	2011	EAI5	EAI5	2010; 2011	Gujarat; Maharashtra, Mumbai	(21, 22)
MCR7	Cattle	Liver	Mumbai, Maharashtra	2012	EAI3_IND	EAI_IND3	2012	Mumbai, Maharashtra / extra-pulmonary	(23)
MCR8	Cattle	Milk	Hyderabad, Telangana	2012	CAS1-Delhi	CAS1_Delhi	2008–2009; 2014–2015; 2010, 2011	Tamil Nadu; Tamil Nadu, Tiruvallur, Southern India; Telangana (formerly Andhra Pradesh) / pulmonary;	(21, 22, 24, 25)
MCR9	Cattle	Nasal swab	Hyderabad, Telangana	2011	T1	T1 T1–T2	2011; 2007–2011 2010; 2014–2015	Tamil Nadu; Bhopal, Madhya Pradesh Hyderabad Telangana (formerly Andhra Pradesh); Southern India	(21, 22, 25, 26)

EAI: East African Indian

CAS: Central and Middle Eastern Asia

U: Unidentified

T1: Tuscany

- a) Sample identification numbers MCR1 and MCR2 were *Mycobacterium tuberculosis* (MT) isolates from wild female ruminant species reported during a previous study (13) and the spoligotyping results of the same were reported during this study
- b) Sample identification numbers MCR3, MCR5, MCR6, MCR7, MCR8 and MCR9 were MT isolates from female cows apart from MCR4 which was from a male; these isolates were identified spoligotyped during this study. All isolates were recovered from cultures in the BACTEC™ MGIT™ 960 system (Beckton Dickinson [BD], Wokingham, United Kingdom), BD-DIFCO™ Middlebrook 7H10 supplemented with oleic albumin dextrose catalase (OADC) and glycerol, and Lowenstein Jensen medium supplemented with glycerol but not sodium pyruvate, and were identified as members of the *Mycobacterium tuberculosis complex* (MTC) group by IS6110 commercial nested polymerase chain reaction (PCR) (GeNei, Bangalore, India). All isolates were confirmed as MT by a specific pattern of reverse slot blot hybridisation of the line probe assay using Genotype® MTC kit (Hain Lifescience, Nehrem, Germany)
- c) The spoligotype patterns were identified among MT isolates; clade designations were made according to SpolDB3
- d) Regional presence of spoligotypes of *Mycobacterium tuberculosis* human isolates (28) are as mentioned in the text (21, 22, 24, 25, 26, 27, 28, 29)