

## CHAPTER 6.5.

# BIOSECURITY PROCEDURES IN POULTRY PRODUCTION

### Article 6.5.1.

#### Introduction

Infectious agents of *poultry* are a threat to *poultry* health and, at times, human health and have significant social and economic implications. In *poultry* production, especially under intensive conditions, prevention is the most viable and economically feasible approach to the control of infectious agents.

*Biosecurity* procedures should be implemented with the objective of preventing the introduction and dissemination of infectious agents in the *poultry* production chain. *Biosecurity* will be enhanced with the adoption and implementation of the principles of Good Agricultural Practices and the Hazard Analysis Critical Control Point (HACCP) system.

### Article 6.5.2.

#### Purpose and scope

This chapter deals with *biosecurity* procedures in intensive *poultry* production. It should be read in conjunction with the Codex Alimentarius Code of Hygienic Practice for Meat (CAC/RCP 58-2005), Code of Hygienic Practice for Eggs and Egg Products (CAC/RCP 15-1976) and Guidelines for the control of *Campylobacter* and *Salmonella* in chicken meat (CAC/GL 78-2011).

This chapter identifies several *biosecurity* measures. The choice of measures to be implemented will vary in accordance with national conditions, including *poultry infection* status, the risk of introduction and dissemination of infectious agents and the cost effectiveness of control measures.

Recommendations on specific infectious agents may be found in relevant disease chapters in the *Terrestrial Code*.

### Article 6.5.3.

#### Definitions

For the purposes of this chapter the following definitions apply:

**Breeders** means *poultry* destined for the production of fertile eggs for incubation for the purpose of producing *day-old birds*.

**Live bird markets** means markets where live birds from various sources and species are sold for *slaughter*, further rearing or production.

Article 6.5.4.

**Recommendations on the location and construction of poultry establishments**

1. All establishments (poultry farms and hatcheries)

- a) A suitably isolated geographical location is recommended. Factors to consider include the location of other *poultry* and livestock *establishments*, *wild* bird concentrations and the distance from roads used to transport *poultry*.
- b) *Poultry establishments* should be located and constructed to provide adequate drainage for the site. Run-off or untreated site waste water should not discharge into waterfowl habitats.
- c) *Poultry* houses and hatcheries should be designed and constructed (preferably of smooth impervious materials) so that cleaning and *disinfection* can be carried out effectively. Ideally, the area immediately surrounding the *poultry* houses and hatcheries should be paved with concrete or other impervious material to facilitate cleaning and *disinfection*.
- d) The *establishment* should be surrounded by a security fence to prevent the entry of unwanted animals and people.
- e) A sign indicating restricted entry should be posted at the entrance to the *establishment*.

2. Additional measures for poultry farms

- a) *Establishments* should be designed to house a single species and a single production type. The design should also consider the “all-in all-out” single age group principle. If this is not feasible, the *establishment* should be designed so that each *flock* can be managed as a separate *epidemiological unit*.
- b) *Poultry* houses, and buildings used to store *feed*, eggs or other material, should be constructed and maintained to prevent the entry of *wild* birds, rodents and arthropods.
- c) Where feasible, the floors of *poultry* houses should be constructed using concrete or other impervious materials and designed so that cleaning and *disinfection* can be carried out effectively.
- d) Where feasible, *feed* should be delivered into the farm from outside the security fence.

3. Additional measures for hatcheries

- a) The design of the hatchery should take account of work flow and air circulation needs, with “one way flow” movement of eggs and *day-old birds* and one way air flow in the same direction.
- b) The hatchery buildings should include physical separation of areas used for the following:
  - i) personnel changing, showering and sanitary facilities;
  - ii) receipt, storage and transfer of eggs;
  - iii) incubation;
  - iv) hatching;
  - v) sorting, sexing and other handling of *day-old birds*;
  - vi) storage of egg boxes and boxes for *day-old birds*, egg flats, chick box liners, chemicals and other items;
  - vii) equipment washing;
  - viii) waste disposal;
  - ix) dining facilities for personnel;
  - x) office space.

Article 6.5.5.

**Recommendations applicable to the operation of poultry establishments**

1. All establishments (poultry farms and hatcheries)

- a) All *establishments* should have a written *biosecurity plan*. Personnel in the *establishments* should have access to basic training in *biosecurity* relevant to *poultry* production and understand the implications to animal health, human health and food safety.
- b) There should be good communication between personnel involved in the *poultry* production chain to ensure that steps are taken to minimise the introduction and dissemination of infectious agents.
- c) Traceability at all levels of the *poultry* production chain should be possible.
- d) Records should be maintained on an individual *flock* basis and include data on bird health, production, medications, *vaccination*, mortality and *surveillance*. In hatcheries, records should include data on fertility, hatchability, *vaccination* and treatments. Records should be maintained on cleaning and *disinfection* of farm and hatchery buildings and equipment. Records should be readily available for inspection on site.
- e) Monitoring of *poultry* health on the *establishment* should be under the supervision of a *veterinarian*.
- f) To avoid the development of antimicrobial resistance, *antimicrobial agents* should be used in accordance with relevant directions of the *Veterinary Services* and manufacturer's instructions and in accordance with Chapters 6.8., 6.9., 6.10. and 6.11.
- g) *Establishments* should be free from unwanted vegetation and debris that could attract or harbour pests.
- h) Procedures for the prevention of entry of *wild* birds into *poultry* houses and buildings, and the control of vermin such as rodents and arthropods should be implemented.
- i) Access to the *establishment* should be controlled to ensure only authorised persons and *vehicles* enter the site.
- j) All personnel and visitors entering an *establishment* should follow a *biosecurity* procedure. The preferred procedure is for visitors and personnel entering the *establishment* to shower and change into clean clothes and footwear provided by the *establishment*. Where this is not practical, clean outer garments (coveralls or overalls, head covering and footwear) should be provided. Entry of visitors and *vehicles* should be registered by the *establishment*.
- k) Personnel and visitors should not have had recent contact with other *poultry*, *poultry* waste, or *poultry* processing plant(s). This time period should be based on the level of risk of transmission of infectious agents. This will depend on the *poultry* production purpose, *biosecurity* procedures and *infection* status.
- l) Any *vehicle* entering an *establishment* should be cleaned and disinfected in accordance with a *biosecurity plan*. Delivery *vehicles* should be cleaned, and disinfected before *loading* each consignment of eggs or *poultry*.

2. Additional measures for all poultry farms

- a) Whenever possible, the "all-in all-out" single age group principle should be used. If this is not feasible and several *flocks* are maintained on one *establishment*, each *flock* should be managed as a separate *epidemiological unit*.
- b) All personnel and visitors entering a *poultry* house should wash their hands with soap and water or sanitize them using a disinfectant. Personnel and visitors should also change footwear, use a boot spray or use a properly maintained disinfectant footbath. The disinfectant solution in the footbath should be changed on a regular basis to ensure its efficacy, in accordance with the manufacturer's instructions.
- c) Any equipment should be cleaned and sanitized before being taken into a *poultry* house.
- d) Animals, other than *poultry* of the appropriate (resident) species and age, should not be permitted access to *poultry* houses. No animals should have access to other buildings, such as those used to store *feed*, eggs or other material.
- e) The drinking water supply to *poultry* houses should be potable in accordance with the World Health Organization or to the relevant national standard, and microbiological quality should be monitored if there is any reason to suspect contamination. The water delivery system should be cleaned and disinfected between *flocks* when the *poultry* house is empty.
- f) Birds used to stock a *poultry* house should preferably be obtained from breeder *flocks* and hatcheries that are free from vertically transmitted infectious agents.

- g) Heat treated *feed* with or without the addition of other bactericidal or bacteriostatic treatments, such as addition of organic acids, are recommended. Where heat treatment is not possible, the use of bacteriostatic or bactericidal treatments is recommended.  
*Feed* should be stored in a manner to prevent access by *wild* birds and rodents. Spilled *feed* should be cleaned up immediately to remove attractants for *wild* birds and rodents. The movement of *feed* between *flocks* should be avoided.
- h) The litter in the *poultry* house should be kept dry and in good condition.
- i) Dead birds should be removed from *poultry* houses as quickly as possible but at least daily. These should be disposed of in a safe and effective manner.
- j) Personnel involved in the catching of birds should be adequately trained in bird handling and basic *biosecurity* procedures.
- k) To minimise stress *poultry* should be transported in well ventilated *containers* and should not be over crowded. Exposure to extreme temperatures should be avoided.
- l) *Containers* should be cleaned and disinfected between each use, or disposed of in a safe manner.
- m) When a *poultry* house is depopulated, it is recommended that all faeces and litter be removed from the house and disposed of in a safe manner to minimise the risk of dissemination of infectious agents.

If litter is not removed and replaced between *flocks* then the litter should be treated in a manner to minimise the risk of dissemination of infectious agents from one *flock* to the next.

After removal of faeces and litter, cleaning and *disinfection* of the *poultry* house and equipment should be done in accordance with Chapter 4.14.

- n) For *poultry flocks* that are allowed to range outdoors, feeders, *feed* and other items which may attract *wild* birds should be kept indoors. *Poultry* should not be allowed access to sources of contamination, such as household waste, litter storage areas, other animals, stagnant water and water of unknown quality. The nesting area should be inside the *poultry* house.

### 3. Additional measures for layers

Refer to Section 3 of the Codex Alimentarius Code of Hygienic Practice for Eggs and Egg Products (CAC/RCP 15-1976).

### 4. Additional measures for breeders

- a) Nest box litter and liners should be kept clean.
- b) *Hatching eggs* should be collected at frequent intervals, at least daily, and placed in new or clean and disinfected packaging materials.
- c) Grossly dirty, cracked, broken, or leaking eggs should be collected separately and should not be used as *hatching eggs*.
- d) *Hatching eggs* should be cleaned and sanitized as soon as possible after collection using an approved sanitising agent, in accordance with the manufacturer's instructions.
- e) *Hatching eggs* or their packaging materials should be marked to assist traceability and veterinary investigations.
- f) The *hatching eggs* should be stored in a dedicated room as soon as possible after cleaning and sanitisation. Storage conditions should minimise the potential for microbial contamination and growth and ensure maximum hatchability. The room should be well ventilated, kept clean, and regularly disinfected using disinfectants approved for this purpose.

### 5. Additional measures for hatcheries

- a) Dead in shell embryos should be removed from hatcheries as soon as they are found and disposed of in a safe and effective manner.
- b) All hatchery waste, garbage and discarded equipment should be contained or at least covered while on site and removed from the hatchery and its environs as soon as possible.
- c) After use, hatchery equipment, tables and surfaces should be promptly and thoroughly cleaned and disinfected with an approved disinfectant.

- d) Egg handlers and sexers and handlers of *day-old birds* should wash their hands with soap and water before commencing work and between working with batches of *hatching eggs* or *day-old birds* from different breeder *flocks*.
- e) *Hatching eggs* and *day-old birds* from different breeder *flocks* should be identifiable during incubation, hatching, sorting and transportation.
- f) *Day-old birds* should be delivered to the farm in new *containers* or in clean, disinfected *containers*.

Article 6.5.6.

**Prevention of further dissemination of infectious agents of poultry**

When a *flock* is suspected or known to be infected, a *veterinarian* should be consulted immediately and, in addition to the general biosecurity measures described previously, management procedures should be adjusted to effectively isolate it from other *flocks* on the *establishment* and other epidemiologically related *establishments*. The following measures are recommended:

- 1) Personnel should manage *flocks* to minimise the risk of dissemination of infectious agents to other *flocks* and *establishments*, and to humans. Relevant measures include handling of an infected *flock* separately, last in sequence and the use of dedicated personnel, clothing and equipment.
- 2) When *infection* has been confirmed, epidemiological investigations should be carried out to determine the origin and route of transmission of the infectious agent.
- 3) *Poultry* carcasses, litter, faeces and other potentially contaminated farm waste should be disposed of in a safe manner to minimise the risk of dissemination of infectious agents. The disposal method used will depend on the infectious agent involved.
- 4) Depending on the epidemiology of the disease, the results of a *risk assessment*, and public and animal health policies, destruction or *slaughter* of a *flock* before the end of the normal production period may be used. When infected *flocks* are destroyed or slaughtered, they should be processed in a manner to minimise exposure of humans and other *flocks* to the infectious agent, and in accordance with recommendations of the *Veterinary Service* and relevant chapters in the *Terrestrial Code*. Based on *risk assessment*, non-infected, high risk *flocks* may be destroyed or slaughtered before the end of their normal production period.

Before restocking, the *poultry* house including equipment should be cleaned, disinfected and tested to verify that the cleaning has been effective. Special attention should be paid to *feed* equipment and water systems.

Microbiological monitoring of the efficacy of *disinfection* procedures is recommended when pathogenic agents have been detected in the previous *flock*.

- 5) Depending on the epidemiology of the disease, *risk assessment*, vaccine availability and public and animal health policies, *vaccination* is an option to minimise the dissemination of the infectious agent. When used, vaccines should be administered in accordance with the directions of the *Veterinary Services* and the manufacturer's instructions. Recommendations in the *Terrestrial Manual* should be followed as appropriate.

Article 6.5.7.

**Recommendations to prevent the dissemination of infectious agents to and from live bird markets**

- 1) Personnel should be educated on the significance of infectious agents and the need to apply *biosecurity* practices to prevent dissemination of these agents. Education should be targeted to personnel at all levels of operations in these markets, such as drivers, owners, handlers and processors. Programmes should be implemented to raise consumer awareness about the risks associated with activities of live bird markets.
- 2) Personnel should wash their hands with soap and water before and after handling birds.
- 3) Birds from diseased *flocks* should not be transported to live bird markets.
- 4) All *containers* and *vehicles* should be cleaned and disinfected every time they leave the market.
- 5) Live birds that leave the market and go to a farm should be kept separately from other birds for a period of time to minimise the potential dissemination of infectious agents of *poultry*.
- 6) Periodically the market should be emptied, cleaned and disinfected. This is of particular importance when an infectious agent of *poultry* deemed significant by the *Veterinary Services* has been identified in the market or the region.

- 7) Where feasible, *surveillance* should be carried out in these markets to detect infectious agents of *poultry*. The *surveillance* programme should be determined by the *Veterinary Services*, and in accordance with recommendations in relevant chapters of the *Terrestrial Code*.
- 8) Efforts should be made to ensure the possibility of tracing all birds entering and leaving the markets.

---

NB: FIRST ADOPTED IN 1982; MOST RECENT UPDATE ADOPTED IN 2013.