

Annex to the Guidance for Completing the OIE template for the collection of data on Antimicrobial Agents intended for use in Animals:

Considerations on converting content of antimicrobial active ingredients in veterinary medicines into kilograms

Calculating the quantities to report in kilogram (kg)

Data on antimicrobial agents intended for use in animals comes in different forms. The OIE template for the collection of data on antimicrobial agents used in animals (OIE template) is designed to collect data on the amounts of chemical compound as declared on the product label. The information may vary, ranging from bulk quantities of antimicrobial agents to numbers of packs of a veterinary medicinal product. The content of antimicrobial agents in such products can be stated in a number of possible ways. It will be necessary, where appropriate, to calculate the required data to populate the OIE template.

Detailed instructions are provided to harmonise some aspects of data reporting:

- Transformation of bulk quantities ([section 1](#)); use this section if you need to convert quantities of raw material, e.g. from import data into the required format.
- Data on veterinary medicinal products ([section 2](#)), including conversion from International Units (IU) to kg (section 2. (ii))
- Recommendations are made in [section 3](#) for further optional conversions, aimed at achieving refined reporting of active entities, the ultimately desired format. If such calculations are made, they should be reported in the OIE template in the free text field provided on the sheets for Reporting Option 1, 2 and 3.

The following abbreviations and symbols will be used:

Symbol/abbreviation	Explanation
Strength	amount of antimicrobial agent per unit of veterinary product
% w/v	per cent weight per volume
mg	milligram
g	gram
kg	kilogram
t	ton (metric)
ml	millilitre
l	litre

1. For data on bulk quantities

Such information is usually sourced from customs, import or other bulk trading. It will likely come as a weight in a number of possible units (e.g. metric tons) of chemical compound and needs to be converted to kg. When conversion into kg is necessary, follow the steps below. If additional conversion factors are needed, please contact the OIE at antimicrobialuse@oie.int.

Step 1: Multiply the amount of antimicrobial agent, i.e. the chemical compound as declared on the product label with the appropriate conversion factor from the table 1 below.

$$\text{Antimicrobial agent (kg)} = \text{antimicrobial agent (unit Z)} \times \text{conversion factor}$$

Table 1: Converting weight units into kg

Unit reported (unit Z)	Conversion factor to kg (for multiplication)
Metric ton	1000
Imperial ton (long)	1016
Imperial ton (short)	907.18
Stone (Imperial)	6.35
Imperial Pound	0.4536
Ounce	0.0283

2. For data on veterinary medicinal products

For veterinary medicinal products containing antimicrobial agents, data on quantities sold is likely to be available as numbers of packages of product sold, with each package containing a specified quantity of medicinal product with a specified amount of antimicrobial agent. In such cases, the amount of antimicrobial agent (chemical compound as declared on the product label) per package needs to be calculated first, and subsequently the result needs to be multiplied with the number of packages of the presentation sold to obtain the overall amount of antimicrobial agent, which should be reported in kg.

The most common ways to indicate the content of the antimicrobial agent(s) of a veterinary medicinal product are:

- (i) Strength in mg or g of the active ingredient per volume or weight or other unit, (for example: ml, l, kg, tablet),
- (ii) Strength in International Units (IU) per weight, volume or other unit,
- (iii) Strength in per cent (%) weight per weight (w/w) or weight per volume (w/v).

Each situation requires a different kind of mathematical conversion.

2. (i) – content of antimicrobial active ingredient (antimicrobial agent) stated in milligram per volume or weight or other unit (for example millilitre, litre, kilogram, tablet) of content

Step 1: Calculation of the content of antimicrobial agent per package

Multiply the amount of antimicrobial agent (chemical compound as declared on the product label) per unit of content, that is, the strength of the product, with the total number of units contained in the package

$$\begin{aligned} & \text{Content of antimicrobial agent per package} \\ & = \text{Strength (amount antimicrobial agent per unit)} \times \text{number of units per package} \end{aligned}$$

Example A:

Tiamulin 100 g/kg premix for medicated feeding stuff; package sizes: (a) 1 kg, (b) 5 kg and (c) 20 kg

Calculation of content of antimicrobial agent, tiamulin, per package:

$$\begin{aligned} (a) \text{ Pack content} &= 100 \text{ g/kg} \times 1 \text{ kg} = 100 \text{ g} \\ (b) \text{ Pack content} &= 100 \text{ g/kg} \times 5 \text{ kg} = 500 \text{ g} \\ (c) \text{ Pack content} &= 100 \text{ g/kg} \times 20 \text{ kg} = 2000 \text{ g} \end{aligned}$$

Example B:

Tetracycline intrauterine tablet containing 2000 mg tetracycline hydrochloride per tablet; package sizes: (a) carton with 1 blister of 5 intrauterine tablets, (b) carton with 4 blisters of 5 intrauterine tablets each (20 tablets), (c) carton with 20 blisters of 5 intrauterine tablets each (100 tablets).

Calculation of content of antimicrobial agent, tetracycline, per package:

$$\begin{aligned} (a) \text{ Pack content} &= 2000 \text{ mg} \times 5 = 2 \text{ g} \times 5 = 10 \text{ g} \\ (b) \text{ Pack content} &= 2000 \text{ mg} \times 20 = 2 \text{ g} \times 20 = 40 \text{ g} \\ (c) \text{ Pack content} &= 2000 \text{ mg} \times 100 = 2 \text{ g} \times 100 = 200 \text{ g} \end{aligned}$$

Example C:

Tilmicosin 300 mg/ml solution for injection for cattle; package sizes: containers of 100 ml and 250 ml; packs of (a) 6, (b) 10 and (c) 12 units of 100 ml and 250 ml.

Calculation of content of antimicrobial agent, tilmicosin, per package:

$$\begin{aligned}
 & \text{(a) Container content} = 300 \text{ mg/ml} \times 100 \text{ ml} = 30000 \text{ mg} = 30 \text{ g} \\
 & \text{Pack content: } \begin{aligned} & \text{(a) } 6 \times 30 \text{ g} = 180 \text{ g,} \\ & \text{(b) } 10 \times 30 \text{ g} = 300 \text{ g} \\ & \text{(c) } 12 \times 30 \text{ g} = 360 \text{ g} \end{aligned} \\
 & \text{(b) Container content} = 300 \text{ mg/ml} \times 250 \text{ ml} = 75000 \text{ mg} = 75 \text{ g} \\
 & \text{Pack content: } \begin{aligned} & \text{(a) } 6 \times 75 \text{ g} = 450 \text{ g,} \\ & \text{(b) } 10 \times 75 \text{ g} = 750 \text{ g} \\ & \text{(c) } 12 \times 75 \text{ g} = 900 \text{ g} \end{aligned}
 \end{aligned}$$

Step 2: Sum up the antimicrobial agent contained in all presentations and packages sold

Convert all contents of antimicrobial agent calculated under step 1 to the same weight unit and add up the total

Step 3: If necessary: convert the total sum of antimicrobial agent contained in all packages of all presentations sold to kg

Multiply the result from step 2 with an appropriate conversion factor to achieve the result in kg

2. (ii) – content of antimicrobial agent (chemical compound as declared on the product label) in International Units (IU) per weight, volume or other unit (for example millilitre, litre, kilogram, tablet) of content

Where the strength of the antimicrobial agent in the veterinary medicinal product is stated International Units (IU) per unit of finished product, an additional conversion step is necessary to obtain results in mg, g, or kg. Table 2 is used to convert content of antimicrobial agents declared in IU on the product label into mg for reporting to the OIE: either divide the total number of IUs of an antimicrobial agent by the value in the column 'International Units (IU) per mg' for this agent in table 2, or, if multiplication is preferred, multiply the total number of IUs with the conversion factor listed for the agent. To convert mg values into kg, please multiply the result of the conversion with 1×10^{-6} equalling 0.000001.

For some antimicrobial agents in veterinary medicinal products, the IU content or strength may be stated in respect to the active entity rather than to the chemical compound actually included; for example: a product may contain penethamate hydroiodide, or procaine benzylpenicillin, but the stated strength in IU refers to benzylpenicillin (product X containing penethamate hydroiodide, equivalent to xx IU benzylpenicillin, or, product Y containing procaine benzylpenicillin, equivalent to yy IU benzylpenicillin). For such cases, use the conversion factor for the relevant active entity listed in table 2 (in the examples used: benzylpenicillin). To convert mg values into kg, please multiply the result of the conversion with 1×10^{-6} equalling 0.000001.

If additional conversion factors are needed or have been used, please contact the OIE at antimicrobialuse@oie.int.

Step 1: Calculating the content of antimicrobial agent per package in IU

Multiply the amount of IU antimicrobial agent per unit of content with the total number of units contained in the package

$$\begin{aligned}
 & \text{Content of antimicrobial agent per package in IU} \\
 & = \text{Strength (amount IU antimicrobial agent per unit)} \times \text{number of units per package}
 \end{aligned}$$

Step 2: Converting the content of antimicrobial agent per package in IU into mg

$$\begin{aligned}
 & \text{Content of antimicrobial agent per package in mg} \\
 & = \text{Content of antimicrobial agent in IU} \times \text{conversion factor}
 \end{aligned}$$

Steps 3-4: Follow steps 2-3 described for (i)

Table 2: Conversion of International Units (IUs) of certain antimicrobial agents into mg and relevant active entities, based on the ESVAC conversion factors¹

Antimicrobial agent in the veterinary medicine	Antimicrobial active entity for reporting to OIE	International Units per mg	Conversion factor to mg for multiplication
Apramycin	Apramycin	556	0.0018
Bacitracin	Bacitracin	74	0.013514
Benzylpenicillin (penicillin G) ²	Benzylpenicillin	1666.67	0.0006
Chlortetracycline	Chlortetracycline	900	0.001111
Colistin methane sulfonate sodium (colistimethate sodium INN)	Colistin	12700	0.000079
Colistin sulfate	Colistin	20500	0.000049
Dihydrostreptomycin	Dihydrostreptomycin	820	0.00122
Erythromycin	Erythromycin	920	0.001087
Gentamicin	Gentamicin	620	0.001613
Kanamycin	Kanamycin	796	0.001256
Neomycin	Neomycin	755	0.001325
Neomycin B (Framycetin)	Neomycin B (Framycetin)	670	0.001492
Oxytetracycline	Oxytetracycline	870	0.001149
Paromomycin	Paromomycin	675	0.001481
Polymyxin B	Polymyxin B	8403	0.000119
Rifamycin	Rifamycin	887	0.001127
Spiramycin	Spiramycin	3200	0.000313
Streptomycin	Streptomycin	785	0.001274
Tetracycline	Tetracycline	982	0.00102
Tobramycin	Tobramycin	875	0.001143
Tylosin	Tylosin	1000	0.001

2. (iii) – content of antimicrobial agent (chemical compound as declared on the product label) in per cent (%) weight per weight (w/w) or weight per volume (w/v) of content

The amount of antimicrobial agent contained in a veterinary medicine concerned may be stated in per cent weight per weight (% w/w) (example 1: product X contains tylosin 100% w/w or, example 2, product Y contains amoxicillin 22.2 % w/w) or in per cent weight per volume (% w/v) (example: product Z contains procaine benzylpenicillin 30% w/v). Such figures first need to be converted into mg/g, g/g, or mg/ml, followed by the calculations described under (i).

Converting % w/w: Conversion calculations are performed by relating the content of antimicrobial agent to 1 g of the finished product. Divide the percentage value by 100 to obtain the amount of antimicrobial agent in g per g finished product.

$$\text{value antimicrobial agent in g per gram finished product} = \frac{\text{value (\%)} \times \text{g}}{100 \text{ (finished product)}}$$

Example 1: Product X containing 100% w/w tylosin will contain 100/100 x g = 1 g tylosin per g finished product.

Example 2: Product Y containing 22.2% w/w amoxicillin will contain 22.2/100 = 0.222 g amoxicillin per g finished product.

Continue with Steps 1-3 of (i)

Converting % w/v: Conversion is based on the assumption that 1 ml of the products weighs 1000 mg. Multiply the percentage value with 10 to obtain the content in mg/ml.

$$\text{value antimicrobial agent in g per ml finished product} = \frac{\text{value (\%)} \times 10 \times \text{mg}}{1 \text{ ml (finished product)}}$$

¹ http://www.ema.europa.eu/ema/pages/includes/document/open_document.jsp?webContentId=WC500189269

² Applies to all derivatives/compounds of benzylpenicillin

Example: Product Z containing 30% w/v benzylpenicillin will contain (30 x 10 mg)/1ml, equal to 300 mg/ml benzylpencicillin.

Continue with Steps 1-3 of (i)

3. Additional recommendations for further conversions of quantities of antimicrobial agents

For pragmatic reasons the OIE accepts the reporting of antimicrobial agents in amounts of chemical compound as declared on the product label of the veterinary medicinal product. However, OIE Member Countries may wish to carry out further calculations to report amounts of active entity. If such further calculations are carried out, please describe them in the OIE template.

Calculating the total amount expressed in weight of chemical compound as declared on the product label of a veterinary medicinal product into antimicrobial active entity (e.g. salt, ester or prodrug into base

This step may be carried out once the steps described in section 1 or section 2. (i) have been completed.

As an example, for the antimicrobial agent tiamulin that is often available in the form of tiamulin hydrogen fumarate (the chemical compound as declared on the product label), the conversion formula to tiamulin (the active entity) would be:

Salt (including base): Tiamulin hydrogen fumarate MW 609.8

Base: Tiamulin MW 493.7

Conversion factor = MW base/MW salt (including base) = 0.81

Multiply the final result in kg obtained by following steps 1 to 3 with the appropriate conversion factor

$$\begin{aligned} \text{Content of active entity (kg)} \\ &= \text{Content of chemical compound as listed on the label (kg)} \\ &\times \text{conversion factor} \end{aligned}$$

Taking the conversion factors of certain derivatives or compounds used by the European Surveillance of Veterinary Antimicrobial Consumption (ESVAC) program managed by the European Medicines Agency, as a starting point, table 3 lists the suggested conversion factors for relevant derivatives or compounds in order to obtain the corresponding amount of the active entity.

If additional conversion factors are needed or have been used, please contact the OIE at antimicrobialuse@oie.int.

Table 3: Conversion of content stated in mg, g or kg of derivatives/compounds of antimicrobial agents in the veterinary product into corresponding mg, g or kg antimicrobial active entity for reporting to the OIE, based on the ESVAC conversion factors³

Derivate or compound	Active entity	Conversion factor for multiplication
Benethamine benzylpenicillin ⁴	Benzylpenicillin	0.61
Benzathine benzylpenicillin	Benzylpenicillin	0.74
Cefapirin benzathine ⁵	Cefapirin	0.78
Cefalexin benzathine ⁶	Cefalexin	0.74
Cloxacillin benzathine ⁷	Cloxacillin	0.78
Oxacillin benzathine ⁸	Oxacillin	0.77
Penethamate hydriodide ⁹	Benzylpenicillin	0.60
Procaine benzylpenicillin ¹⁰	Benzylpenicillin	0.57

Step 1–3: As described in section 2. (i)

Step 4: Multiply the final result in kg obtained by following steps 1 to 3 with the appropriate conversion factor listed in table 3

$$\begin{aligned}
 & \textit{Antimicrobial agent (active entity)(kg)} \\
 & = \textit{antimicrobial agent (chemical compound as declared on the product label)(kg)} \\
 & \quad \times \textit{derivate or compound conversion factor}
 \end{aligned}$$

³ http://www.ema.europa.eu/ema/pages/includes/document/open_document.jsp?webContentId=WC500189269

⁴ Conversion factor for benethamine benzylpenicillin is updated from 0.65 to 0.61

⁵ Conversion factor for cefapirin benzathine is updated from 0.41 to 0.78

⁶ Conversion factor for cefalexin benzathine is updated from 0.36 to 0.74

⁷ Conversion factor for cloxacillin benzathine is updated from 0.43 to 0.78

⁸ Conversion factor for oxacillin benzathine is updated from 0.69 to 0.77

⁹ Conversion factor for penethamate hydriodide is updated from 0.63 to 0.60

¹⁰ Conversion factor for procaine benzylpenicillin is updated from 0.61 to 0.57